



Product:

Zetadex-50 Medium

Gel Filtration Matrix

Product Code:	TM-0106
CAS Number:	9048-71-9
HTS Code:	39139000990

Application Note

Zetadex-50 Medium is a size-exclusion matrix. Molecules purified with Zetadex-50 Medium are separated according to size. Smaller molecules pass significantly slower through the column than larger molecules. Buffer and pH effects on resolution are minimal. The size exclusion cut-off for Zetadex-50 Medium is set at 25 kD for proteins and 20 bp for nucleic acids. Purified biomolecules are not significantly diluted when processed using Zetadex-50 Medium.

Hydration, Filling and Packing Columns

Zetadex-50 Medium can be hydrated in aqueous media of choice containing no more than 20% alcohol. Time of hydration should be a minimum of 3 hours at room temperature or 1 hour at 90°C. Hydrated Zetadex-50 Medium is provided in a settled gel volume to buffer volume ratio of 1:1, degassed and ready to use. Filling a column requires that the slurry be not too thick as to retain air bubbles. A settled gel volume to buffer volume ratio of 3:1 is optimal for this purpose. The remaining buffer may be used later for column packing.

To fill, pour the Z-50 slurry into a tilted column. Alternatively, a funnel may be used, with the tip of the funnel touching the inside wall of the column to prevent splashing of the slurry. A gel reservoir or column extension is desirable for filling the whole column in a single operation. To finish packing the gel bed, a peristaltic pump may be employed. Use as high a flow rate as possible without deforming the beads. Zetadex-50 Medium can be pressurized up to 3 bar.

Precautions for Safe Handling

As part of good industrial and personal hygiene and safety procedure, avoid all unnecessary exposure to the chemical substance and ensure prompt removal from skin, eyes and clothing. Keep container tightly closed. Suitable for any general chemical storage area. Containers of this material may be hazardous when empty since they retain product residues (dust, solids); observe all warnings and precautions listed for the product.

Other Product Properties

Bead structure: Bead size (Dry): Bead size (Wet): Maximum operating pressure:	Cross-linked spherical dextran composite 50-150 µm 100-300 µm Generally obeys Darcy's Law: $U = K_0 \frac{\Delta P}{L}$	
	where U = linear flow rate, cm/hour ΔP = pressure drop over gel bed, cm H ₂ O L = bed ht, in cm K _o = 145 for Z-50 Medium	
Chemical stability:	All commonly used buffers, including: 0.2M NaOH; 0.2M HCl; 1M acetic acid; 8M urea; 6M guanidine HCl; 1% SDS; 24% Ethanol; 30% Propanol; 30% Acetonitrile.	
pH stability:	2.0 to 10.0	
Autoclavable:	121°C, pH 7 for 30 minutes	
Storage & Shipping	Ambient	
	The use of this product is strictly limited to trained personnel for professional manufacturing laboratory or research purposes. Final Fitness-For-Lise must	

The use of this product is strictly limited to trained personnel for professional manufacturing, laboratory, or research purposes. Final Fitness-For-Use must be determined by and is the sole responsibility of the end-user.

Document: CM-06-Instr_v2 Version: CHR-7006-02