

Instructions For Use

Product: **Zetadex-50 Agglutination Grade**
Speciality Zetadex Gel Resin optimized for blood typing using Gel Card Technology

Product Code: TM-0111
 CAS Number: 9048-71-9

Application Note

Zetadex-50 Agglutination Grade is a size-exclusion matrix. Molecules purified with Zetadex-50 Agglutination Grade are separated according to size. Smaller molecules pass significantly slower through the column than larger molecules. Buffer and pH effects on resolution are minimal. The size exclusion cut-off for Zetadex-50 Agglutination Grade is set at 25 kD for proteins and 20 bp for nucleic acids. Purified biomolecules are not significantly diluted when processed using Zetadex-50 Superfine.

Hydration and Filling

Zetadex-50 Agglutination Grade can be hydrated in aqueous media of choice containing no more than 20% alcohol. Time of hydration should be a minimum of 3 hours at room temperature or 1 hour at 90°C. Hydrated Zetadex-50 Agglutination Grade is provided in a settled gel volume to buffer volume ratio of 1:1, degassed and ready to use. Filling requires that the slurry be not too thick as to retain air bubbles. A peristaltic pump may be employed. Use as high a flow rate as possible without deforming the beads. Zetadex-50 Superfine can be pressurized up to 3 bar.

Precautions for Safe Handling

As part of good industrial and personal hygiene and safety procedure, avoid all unnecessary exposure to the chemical substance and ensure prompt removal from skin, eyes and clothing. Keep container tightly closed. Suitable for any general chemical storage area. Containers of this material may be hazardous when empty since they retain product residues (dust, solids); observe all warnings and precautions listed for the product.

Other Product Properties

Bead structure: Cross-linked spherical dextran composite
 Bead size (Dry): 20-50 µm
 Bead size (Wet): 40-100 µm
 Maximum operating pressure: Generally obeys Darcy's Law: $U = K_0 \frac{\Delta P}{L}$
 where U = linear flow rate, cm/hour
 ΔP = pressure drop over gel bed, cm H₂O
 L = bed ht, in cm
 K₀ = 13.5 for Z-50 Superfine

Chemical stability: All commonly used buffers, including: 0.2M NaOH; 0.2M HCl; 1M acetic acid; 8M urea; 6M guanidine HCl; 1% SDS; 24% Ethanol; 30% Propanol; 30% Acetonitrile.
 pH stability: 2.0 to 10.0
 Autoclavable: 121°C, pH 7 for 30 minutes
 Storage & Shipping: Ambient

The use of this product is strictly limited to trained personnel for professional manufacturing, laboratory, or research purposes. Final Fitness-For-Use must be determined by and is the sole responsibility of the end-user.